

Regulation of milk protein solubility by a whey-derived proline-rich peptide product

Sophie N. Selby-Pham^{1,2}, Kate Howell², Hema Jegasothy¹, Paul Sheean¹, Tanoj Singh¹, Cheryl Taylor¹ and Louise E. Bennett^{1*}

¹CSIRO Animal, Food and Health Sciences, 671 Sneydes Road, Werribee, Victoria, 3030, Australia

²Department of Agriculture and Food Systems, Melbourne School of Land and Environment, University of Melbourne, Parkville, Victoria, 3051, Australia

Received 23 August 2012; accepted for publication 8 February 2013; first published online 23 April 2013

The effects of a bovine whey peptide product enriched in proline (wPRP) on the solubility of milk proteins were tested under ambient conditions or following heat treatment at 75 and 100 °C, for 1 and 15 min, followed by post-incubation storage at either ambient temperature or 4 °C for up to 7 d. wPRP promoted solubilisation of milk proteins in a concentration-dependent manner without heat treatment and also after heat treatment at 75 and 100 °C, and the effect was enhanced after storage under either ambient or refrigerated storage conditions. Interactions of wPRP and milk proteins were monitored by particle size analysis and tryptic digestion and specifically linked with solubilisation of α S1 casein (α S1-Cn), which supported observed changes in milk protein solubility. The results suggested that wPRP preferably prevented or reversed physical versus covalent protein aggregation, with the relaxation of hydrophobic interactions at 4 °C providing an additive effect. This application of wPRP represents a novel approach to stabilisation of dairy proteins following thermal processing with industrial usefulness yet to be explored.

Keywords: Protein aggregation, proline-rich peptide, hydrophobic interaction.

Abbreviations: WPI, whey protein isolate; SMP, skim milk powder; wPRP, whey proline rich peptide.

Thermal processing is essential for microbial stabilisation of milk but also induces chemical and physical interactions leading to protein aggregation and precipitation, that may produce either favourable or undesirable functional properties (Singh, 2004). Manipulation of heating conditions continues to represent the key industrial processing parameter used to control the functional properties of milk proteins (Yuksel & Erdem, 2005; Raikos, 2010). The effects of heating conditions including time, temperature, pH, additives that react or interact with proteins, and manipulation of the protein and non-protein fractions of milk, have been extensively studied, with a view to understanding the mechanistic nature of interactions (Relkin & Mulvihill, 1996; Singh, 2004; Donato & Guyomarch, 2009) and importantly, the consequences for dairy product quality (Guyomarc'h et al. 2003; Anema, 2008; Morand et al. 2011).

In skimmed milk, interactions between whey proteins and caseins are accelerated as a function of temperature

and heating time (Anema & Li, 2003) and intermolecular aggregation is observed at and above temperatures that invoke denaturation of whey proteins (>60 °C), (Parris, 1993; Relkin & Mulvihill, 1996; Parris et al. 1997). However, following whey to casein protein cross-linking induced during pre-heating, progressive destabilisation of micellar and non-micellar protein aggregates involving both whey proteins and caseins, can lead to undesirable protein sedimentation and eventually phase separation in ultra-high temperature (UHT)-treated milk (Datta & Deeth, 2001), concentrated milk (Muir & Sweetsur, 1978), homogenised milk (McCrae & Muir, 1992) and reconstituted milk (Chandrapala et al. 2010).

In situations where protein aggregation and precipitation is undesirable, numerous approaches have been trialled to maintain milk protein solubility. One approach has been to manipulate the mineral balance in milk with calcium-chelating agents such as sodium or potassium salts of phosphate (Williams et al. 2005) and citrate (Augustin & Clarke, 1990), so as to maintain stability of the colloidal calcium phosphate (CCP)-casein micelle phase

*For correspondence; e-mail: louise.bennett@csiro.au

(Chandrapala et al. 2010). Alternative approaches for improving heat stability of homogenised milk include the addition of urea (Muir & Sweetsur, 1977), phospholipid-based emulsifiers (McCrae, 1999), such as soy or egg lecithin (Le et al. 2011), polysaccharides such as carrageenan, alginate and gum arabic (Ye, 2008), and products generated by phospholipases increased the heat-stability of β -lactoglobulin (β -lg) in whey protein concentrate (Havn et al. 2006). Stabilising effects against thermal aggregation have also been achieved using phenolic compounds (O'Connell & Fox, 1999a, b).

Isolated whey proteins exhibit particular tendency for thermal instability due to their globular nature. However, the solubility of whey proteins can be maintained during and after thermal processing by removal of salts (de Rham & Chanton, 1984), addition of agents to complex free mineral ions (Anand & Ward, 2005) or by stabilising whey proteins against thermal denaturation (Kulmyrzaev et al. 2000). Other approaches include adopting heating conditions favouring assembly of small aggregates that can be physically redispersed using shear (APV LeanCream™ process, SPX). Thermal stability of whey proteins was also regulated by controlling protein denaturation using addition of minerals optimised to the concentration of total solids, forming a soluble concentrate of whey proteins (Gao et al. 2008). Modification of whey proteins by hydrolysis has also been used to make them suitable for several purposes including heat stability (Foegeding et al. 2002).

Conversely, milk caseins have been applied to regulate aggregation of a wide range of proteins which is attributed to their 'molecular chaperone' properties and capacity to function like small heat shock proteins (Thorn et al. 2005, 2008, 2009). Subsequently, a selection of dairy protein hydrolysates were reported to inhibit self-assembly of both reduced, carboxymethylated kappa casein (RCM- κ -Cn) and human amyloid beta peptide into fibrils, presumably by a similar mechanism (Bennett et al. 2009). We have recently shown that the proline-rich peptide (PRP) product prepared from whey proteins (wPRP), as used in the present study, prevented assembly of beta amyloid, as implicated in Alzheimer's Disease, into either oligomers or fibrils and thereby modulated its toxic effects on cells (Bharadwaj et al. 2012). While the manipulation of ratios of proteins can alter extents of thermal aggregation (Yong & Forgeding, 2010), the usefulness of specific proteins or hydrolysates for assisting in thermal processing of milk and dairy products has yet to be systematically applied and we therefore sought to investigate the efficacy of wPRP in this novel application.

In this study, we report efficacy of wPRP on regulating the solubility of milk proteins either without or with heating at 75 and 100 °C, and subsequent effects of ambient temperature and refrigerated storage. We substantiate effects of wPRP on milk protein solubility by total protein solubility assay, particle size analysis and identify specific proteins affected by wPRP, using capillary electrophoresis. We interpret the effects of wPRP in terms of 'chaperone-like', reversible binding interactions that can inhibit and reverse physical

aggregation processes and thereby promote solubility of milk proteins. The results suggest that this novel approach to dairy protein stabilisation may be useful for industrial applications.

Materials and methods

Materials

Whey protein isolate (WPI, containing 90% protein, 3.7% ash, 0.7% Na, 0.3% P and 0.15% Ca, w/w) was obtained from Murray Goulburn, (Natrpro, MG Nutritionals, Brunswick, Australia). Fresh skimmed milk, purchased from a local supermarket (Woolworths Dairy Fresh Skim Milk, NSW, Australia), was freeze-dried and stored at 4 °C. The skimmed milk powder (SMP) contained 37.1% (w/w) true protein (calculated as total nitrogen minus non-protein nitrogen $\times 6.38$). Citric acid, DL-dithiothreitol (DTT), 2-[N-Morpholino] ethanesulfonic acid (MES), glycyl-L-tyrosine, trypsin (for tryptic digestion), trifluoroacetic acid (TFA, spectrophotometric grade $\geq 99\%$) and urea (Electrophoresis grade) were obtained from Sigma-Aldrich, St. Louis, MO, USA. Tri-sodium citrate (Analar) was obtained from B.D.H. Ltd, Dorset, UK. Acetonitrile and ethanol were from Ajax Fine Chemicals, NSW, Australia and methyl hydroxyethyl cellulose was from TCI Development Co. Ltd, Shanghai, China.

Preparation of whey protein hydrolysate and C18 solid phase extract

Whey protein hydrolysate was prepared from WPI as described previously (Bharadwaj et al. 2012) before isolation of the wPRP product by C18 solid phase extraction as follows. The non-binding peptide mixture (containing 12.2% nitrogen) recovered after preparative ion-exchange chromatography, was dissolved in de-ionised water at 40 mg/ml ('crude' wPRP) before loading onto a pre-conditioned C18 solid phase cartridge (Strata-X 33 μ m Polymeric Reverse Phase cartridges, 500 mg/6 ml, Phenomenex, CA, USA). The cartridges were washed, equilibrated and then loaded with 5.0 ml each of methanol, water and 'crude' wPRP solution, respectively. Non-binding solids were eluted in a further 5.0 ml water and the bound fraction containing wPRP's was eluted with 5.0 ml 100% acetonitrile. The wPRP product was dried by centrifugal evaporator under vacuum (Genevac, Thermo Fisher Scientific, Waltham, MA, USA) and stored at -20 °C.

Milk protein solubility assay

The effects of added wPRP on milk protein solubility following experimental heat treatments were measured as follows. SMP and wPRP were reconstituted in 10 mM MES buffer pH 6.5 (MES buffer), at stock concentrations of 100 and 20 mg/ml, respectively, and incubated at ambient

temperature for 1 h with occasional gentle mixing. SMP and wPRP mixtures, at final concentrations of 10 and 0 to 5 mg/ml, respectively, were prepared in MES buffer and cooled on ice immediately following thermal treatment. Replicates of SMP±wPRP were tested either immediately (ambient temperature) or after 1 or 7 d storage at 4 °C.

Soluble protein and other species remaining soluble after centrifugation (25 000 g at 4 °C for 15 min, Model 5417R, Eppendorf AG, Hamburg, Germany) were measured by absorbance at 280 nm (UV-1650 PC spectrophotometer, Shimadzu, Kyoto, Japan), adapted from the method of Laleye et al. (2008). Supernatants were diluted by 1/2 in 10 mM MES buffer, pH 6.5 before measuring absorbance at 280 nm. The contribution of light scatter to the measured absorbance at 280 nm was corrected using the absorbance at 340 nm, and the relationship that light scatter, at wavelength λ , is proportional to $1/\lambda^4$ (Xu, 2001). 'True' absorbance was thereby calculated from the measured absorbance at 280 nm using Equation 1:

$$A_{280 \text{ true}} = A_{280 \text{ measured}} - \left\{ \frac{340^4}{280^4} \right\} \times A_{340} \quad (1)$$

A calibration profile of unheated wPRP was included in each experiment to account for the contribution of wPRP to the absorbance at 280 nm, over the experimental concentration range of wPRP (0–5 mg/ml). 'True' wPRP (calculated by linear regression) was subtracted from 'True' SMP+wPRP according to Equation 2:

$$A_{280 \text{ true(SMP)}} = A_{280 \text{ true(SMP+wPRP)}} - A_{280 \text{ true(wPRP)}} \quad (2)$$

Standardised changes in SMP protein solubility (%smpsol) were expressed using light scatter and wPRP-corrected data, according to Equation 3:

$$\% \text{smpsol} = \frac{A_{280 \text{ (SMP+wPRP at } T)}}{A_{280 \text{ (SMP at } T)}} - \frac{A_{280 \text{ (SMP at } T)}}{A_{280 \text{ (SMP at } T)}} \times 100 \quad (3)$$

where %smpsol is the change in solubility of SMP proteins in the presence ($A_{280 \text{ (SMP+wPRP at } T)}}$) and absence of wPRP ($A_{280 \text{ (SMP at } T)}}$), treated at temperature T . All data points represent the mean of sample preparations in triplicate.

Particle size analysis

Sample sets of reconstituted SMP at a final concentration of 20 mg/ml in the absence or presence of wPRP at 7 mg/ml and suitable controls, were prepared in MES buffer. The samples were analysed after 15 min (Day 0, stored at 22 °C), 1 d (Day 1, stored at 22 °C) and 15 d (Day 15, stored at 4 °C, no evidence of microbial spoilage was detected in these samples by visual inspection). Samples were placed on ice pending analysis and suitable dilutions of the whole samples were prepared prior to measurement. Particle size analysis was conducted by dynamic light scattering using a Zetasizer Nanoseries Nano-ZS (Malvern, Worcestershire, UK) detecting particles from 0.3 nm to 10 μm diameter at a scattering angle of 173°, with temperature controlled at 25 °C. Results

are expressed as the mean of duplicate acquisitions of at least 5 replicates of each sample.

Tryptic digestion and peptide profiling by reversed phase-HPLC

The Day 0 sample (22 °C for 15 min) prepared for particle size analysis was diluted by 1/2 in MES buffer (10 mg/ml SMP, 3.5 mg/ml wPRP) and digested with trypsin. Trypsin stock solution was freshly prepared in MES buffer and added to samples at a final concentration of 2% (w/w) before incubating at 37 °C for 3 h. Trypsin activity was stopped by addition of TFA to 1% (v/v). Samples were prepared and analysed in triplicate.

The tryptic digests of wPRP and SMP and suitable controls were analysed by C18 reverse phase high performance liquid chromatography (RP-HPLC) using a Vydac 238EV52 monomeric Everest[™] C18 column (250 mm × 2.1 mm, 5 μm particle size, 300 Å pore size with guard column, 10 mm × 2.1 mm, Grace Davidson Discovery Sciences, Rowville, Australia) and automated ThermoFinnigan Surveyor Plus HPLC system (Thermo Fisher Scientific Australia Pty Ltd, Scoresby, Australia). The column temperature was 35 °C and separation was conducted with a linear gradient of 5 to 40% solvent A (0.08% TFA in acetonitrile) in solvent B (0.1% TFA in de-ionised water) over 40 min, then Solvent A was increased linearly to 60% over 10 min and maintained for a further 10 min. The flow rate was constant at 150 $\mu\text{l/min}$ and monitoring was at 214 nm by photodiode array detector (Thermo Fisher Scientific Australia).

Capillary electrophoresis

Selected samples prepared at the same ratio of SMP to wPRP as used for particle size analysis were analysed by capillary electrophoresis (CE), after diluting the complete sample in MES buffer (1/2), then centrifuging (25 000 g, at 4 °C for 15 min) and further diluting the supernatant in MES buffer (1/2). Samples were prepared after 15 min at 22 °C (Day 0) and 24 h at 4 °C (Day 1). Samples and milk protein standards (5 mg/ml protein) were mixed in CE sample buffer (5.0 mM tri-sodium citrate, 9.9 mM DTT and 6.0 M urea, pH 8.0) at 1:5 (total volume 720 μl) plus 40 μl of internal standard (42 mM glycyl-L-tyrosine in de-ionised water) before incubating for 1.5 h at 22 °C and then filtering (0.22 μm , Acrodisc syringe filters, Pall Corporation, Ann Arbor, MI, USA).

Analysis was conducted using a Beckman Coulter P/ACE MDQ CE system (Beckman Coulter, CA, USA) using a 50 μm i.d. uncoated fused silica capillary with an effective length of 30 cm (Beckman Coulter, Fullerton, California, USA) and UV detection at 214 nm, based on the method of Kanning et al. (1993). Prior to sample injection, the capillary was rinsed under pressure at 138 kPa for 1.0 min each with the following sequence of reagents: 0.5% sodium dodecyl sulphate; deionised water; 0.1 M sodium hydroxide; de-ionised water and run buffer (containing 20 mM tri-sodium citrate, 0.5 mg/ml methyl hydroxyethyl cellulose, 6.0 M urea,

adjusted to pH 3.0 with citric acid). Standards and samples were pressure injected for 10 s at 7 kPa, and separated under normal polarity at 20 kV and 35 °C. Peak identifications were made by comparing mobilities with protein standards and referring to Kanning et al. (1993) but substituting reference to α S0 casein (α S0-Cn) for ' α S1-P-Cn', representing the phosphorylated form of α S1-Cn at serine position 41.

Statistical analysis

Analysis of variance was conducted by Dunnett method to determine if the addition of wPRP resulted in a significant change in SMP protein solubility compared with the control (no wPRP) at the same heat treatment. Peak area results of milk proteins obtained from CE analysis were subjected to Tukey's test with $\alpha=5\%$. MiniTab Pro 16.1.0.0. (Minitab Inc., State College, PA, USA) was used for all statistical analyses.

Results and discussion

Effects of wPRP on solubility of milk proteins

The effects of wPRP on solubility of milk proteins as a function of heating temperature and time was measured by the milk protein solubility assay. Non-aggregating proteins dissolved at appropriate concentrations (absorbance at 280 nm ≤ 0.5) do not absorb at 340 nm, as supported by the response shown for wPRP (Fig. 1a). This justifies the previously reported approach (see Methods) of using the absorbance at 340 nm as a means of correcting for (i.e., subtracting) the contribution of light scatter (due to the presence of soluble aggregates), to the total absorbance at 280 nm. The low absorbance at 340 nm after incubation of wPRP alone also indicated that the tendency for self-aggregation of wPRP was negligible at 22 °C (Fig. 1a). Self-aggregation of wPRP alone at elevated temperatures was not considered as wPRP was always tested in the presence of SMP protein in the solubility assay.

In general the presence of wPRP during heating increased milk protein solubility in a concentration-dependent manner to an extent that depended on the heat treatment and the subsequent storage time at 4 °C. For samples held at 22 °C for 15 min (Fig. 1b), the immediate change in milk protein solubility (Day 0) reached approximately 20% at the maximum level of wPRP of 5 mg/ml. After storage at 4 °C for 1 d, the presence of wPRP at 5 mg/ml further increased SMP protein solubility by >63% (Fig. 1b). These effects of wPRP demonstrated the capacity for further dissolution of SMP proteins after the standard sample preparation method. The relationship between Day 0 and Day 1 data (diverging from parallel lines) suggested that the combination of low temperature and the presence of wPRP favoured increased solubility of SMP protein to a greater extent than temperature alone.




Fig. 1. (a) Standard curves for whey proline-rich peptide product (wPRP) monitored at 280 and 340 nm. (b) Effects of wPRP on standardised change in milk protein solubility following treatment at 22 °C for 15 min (Day 0) and of samples analysed after storage for 1 d at 4 °C (Day 1). Data points labelled '*' are significantly different from 0 ($P < 0.05$, ANOVA test, Dunnett Method). Results represent the mean of sample analysis in triplicate and error bars represent standard error of the mean.

Under heating conditions at 75 °C for 1 min, where whey protein denaturation and limited protein-protein cross-linking are expected, the presence of wPRP at 5 mg/ml increased protein solubility to a maximum of 51 and 93% at Day 1 and Day 7, respectively (Fig. 2a). The presence of wPRP at 5 mg/ml was more effective at maintaining SMP protein solubility after treatment for 1 min (Fig. 2a) vs. 15 min, where the maximum increase in protein solubility was 30 and 60% at Day 1 and Day 7, respectively (Fig. 2b). As for studies at 22 °C, SMP protein solubility was significantly improved after incubation at 4 °C for 7 d. wPRP was more effective at preventing loss of solubility of SMP heated for 1 min vs. 15 min at either storage time, suggesting that wPRP was relatively less effective at




Fig. 2. Effects of whey proline-rich peptide product (wPRP) on standardised change in milk protein solubility following treatment at (a) 75 °C for 1 min; (b) 75 °C for 15 min; (c) 100 °C for 1 min and (d) 100 °C for 15 min. Results are shown either immediately after heat treatment (Day 0), after storage at 4 °C for 1 d (Day 1) or 7 d (Day 7), as labelled. The concentration of SMP was 10 mg/ml. Data points labelled '*' are significantly different from 0 ($P < 0.05$, ANOVA test, Dunnett Method). Results represent the mean of sample analysis in triplicate and error bars represent standard error of the mean.

preventing thermo-chemical protein association compared with physical aggregation processes.

At 100 °C and above 2 mg/ml wPRP, SMP protein solubility was improved at both at Day 0 and Day 7 for heating at either 1 min (Fig. 2c) or 15 min (Fig. 2d). Further increase in protein solubility was evident following 7 d storage at 4 °C, with increases in solubility of >75% at 5 mg/ml wPRP (Fig. 2c, d, Day 7). These results further support that wPRP was not as effective in preventing covalent reactions associated with milk protein heating, namely, κ -Cn and β -lg sulphhydryl-disulphide cross-linking (Anema & Li, 2003), isopeptide bond formation and dehydroalanine-mediated cross-links (O'Connell & Fox, 2003), but could disperse physical aggregates during refrigerated storage.

Dispersion of milk protein physical aggregates by wPRP

Particle size analysis was conducted to independently verify the effects of wPRP on aggregation behaviour of SMP proteins as observed in the SMP solubility assay at 22 °C (Fig. 1b). The ratios of wPRP to SMP in samples corresponded to mixtures at 3.5 mg/ml wPRP shown in Fig. 1b. In the absence of wPRP, the mean particle size increased slightly from 169.9 ± 0.6 to 175.5 ± 0.5 nm after 15 d storage at 4 °C (z-average value, Fig. 3a). However, in the presence of wPRP, the mean particle size decreased significantly ($P < 0.05$) from 185.0 ± 0.6 to 140.7 ± 1.8 nm (z-average values, Fig. 3b). This supported that wPRP was able to disperse physical aggregates present in unheated SMP




Fig. 3. Particle size distribution of (a) SMP alone and (b) SMP incubated with whey proline-rich peptide product (wPRP), expressed as changes in particle size relative abundance as a percentage. The concentrations of SMP and wPRP were 20 and 7.0 mg/ml respectively and incubation conditions were: 22 °C for 15 min (Day 0); 22 °C for 24 h (Day 1, 22 °C), followed by 4 °C for 15 d (Day 15). Results represent the mean of duplicate acquisitions of at least 5 replicates of each sample.

and thereby promoted solubility of reconstituted milk proteins.

These results collectively suggested that the presence of wPRP had a positive effect on maintaining or increasing the solubility of milk proteins either at room temperature or following heating at 75 and 100 °C, and that solubility was further enhanced by storage at 4 °C. The changes in solubility that occurred post-storage indicated that wPRP was able to dynamically disperse physical protein aggregates. The weakening of hydrophobic interactions as a function of temperature (Baldwin, 1986) would be expected to increase the solubility of SMP proteins even in the absence of wPRP, however, the divergence from parallel of




Fig. 4. Comparison of amino acid profiles of whey protein isolate substrate and whey proline-rich peptide (wPRP) product. Analysis by Pearson Product Moment Correlation indicated there were no significant relationships between any pair of variables ($P > 0.050$).

solubility curves measured pre- and post-storage (Fig. 2a, c, d), inferred a positive interactive effect of wPRP and low temperature storage.

Proposed role of proline-rich peptides in regulating milk protein solubility

As a result of the method of processing, wPRP was compositionally distinct from the WPI substrate according to its amino acid profile (Fig. 4). The wPRP product was relatively enriched with respect to the hydrophobic amino acids proline and phenylalanine compared with WPI, and the effects of this product on regulating self-assembly of beta amyloid peptide was recently described (Bharadwaj et al. 2012).

PRPs have been previously isolated from caseins rather than whey proteins as β -casein (β -Cn) represents a particularly rich source of proline (35/209 residues or 17%, w/w) compared with 2.5% (w/w) found in WPI. However, the process described for preparation of wPRP was successful in producing a proline-rich peptide fraction from whey, enriched to 100.1 mg/g or 10.0% w/w (Fig. 4) and we hypothesise that the properties of wPRP are strongly linked with the elevated proline content.

The term 'proline-rich peptide' has been used to describe peptides containing a high proportion of proline and hydrophobic amino acids. The level of enrichment of proline, phenylalanine and other hydrophobic amino acids justifies its characterisation as a proline-rich peptide (Janusz et al. 1981). A number of biological effects of PRPs have been reported with potential for nutritional and therapeutic uses but as yet, PRPs do not appear to have been tested or exploited for technological uses such as in processing of milk or other food proteins. Regulation of the assembly of fibrillar structures displayed by specific purified




Fig. 5. Analysis of soluble proteins detected following incubation of SMP (10 mg/ml) in the absence or presence of whey proline-rich peptide product (wPRP, 3.5 mg/ml) following incubation at 22 °C for 15 min (Day 0) or 4 °C for 24 h at (Day 1, 4 °C), by capillary electrophoresis. (a) representative electropherograms from triplicate analysis. (b) Mean peak areas of proteins standard detectable without interference from wPRP's at Day 0 and Day 1. Results represent the mean of sample analysis in triplicate after correction of peak areas for the internal standard and error bars represent standard error of the mean. Significant effects of wPRP on enhancing solubility are shown (*, $P < 0.05$, Tukey's test). α La: α -lactalbumin; β Lg: β -lactoglobulin; α S1-Cn: α _{S1}-casein; α S1-P-Cn: α _{S1}-casein with serine 41 phosphorylated; κ -Cn: κ -casein; β -Cn-A1: β -casein A1; β -Cn-A2.

milk casein proteins by other specific milk caseins shows the potential for co-operative interactions between this particular group of proteins. For example, the studies of Thorn et al showed that in purified form, while the caseins: κ -Cn and α S2 casein (α S2-Cn) can self-assemble into β -sheet-rich fibrillar structures, in contrast, β -Cn and α S1-Cn prevent formation of these fibrillar structures (Thorn et al. 2005, 2008, 2009). Furthermore, we have recently shown that hydrolysates prepared from either whey




Fig. 6. Representative HPLC profiles of tryptic digests of SMP alone, SMP incubated with whey proline-rich peptide product (wPRP) and wPRP alone, following incubation of SMP (10 mg/ml) at 22 °C for 15 min ('Day 0') in the absence or presence of wPRP (3.5 mg/ml).

or casein proteins can also inhibit assembly of fibrils of both RCM- κ -Cn and amyloid beta peptide, associated with Alzheimer's Disease (Bharadwaj et al. 2012). The previous authors use the term 'molecular chaperone' to describe these phenomena, and we have likewise adopted this descriptive term even though the ratios of substrate to 'chaperone' are not strictly analogous to true chaperone-substrate interactions.

'Colostrinin' (CLN) is a mixture of low molecular weight PRPs derived from ovine colostrum (Zimecki, 2008), containing non-polar amino acids and proline contents of 40 and 25%, respectively. The extensive research characterising properties of CLN suggests that the proline-rich nature of this peptide extract is important in its functional and bioactive properties (Janusz & Zablocka, 2010). In addition to biological effects, the effects of CLN on structure of the fibrillogenic peptide in Alzheimer's disease, beta amyloid 1–42 (A β 42), have been demonstrated. It has been reported that CLN not only inhibited the aggregation of A β 42 into cross β -sheet-rich fibrils but also dissolved pre-existing fibrils of A β 42 in vitro (Schuster et al. 2005). Likewise, we have recently shown that wPRP can regulate oligomer and fibril assembly of A β 42 (Bharadwaj et al. 2012), and propose that the prevention of physical aggregation of milk proteins by wPRP is related to effects exerted on A β 42. We suggest that hydrophobic binding interactions between wPRP and milk proteins, specifically α S caseins (Fig. 5a, b), and storage effects similarly reflect progressive disruption of α - and β -sheet secondary structures, as shown for CLN (Boldogh et al. 2008).

Specific interactions of wPRP and milk proteins

Replicates of samples prepared as for particle size analysis were subjected to digestion with trypsin in order to determine if binding interactions between wPRP and milk proteins were strong enough to alter accessibility to tryptic

cleavage sites. After accounting for the presence of wPRP itself, there were at least 4 peaks that appeared to be altered by the presence of wPRP: 3 with decreased abundance and 1 with increased abundance (Fig. 6). Considering the high number of unaffected peptides, this suggested that interactions between wPRP and milk proteins exerted only limited interference to hydrolysis by trypsin. This is supported by the fast dissociation kinetics reported to take place between proline-rich domains of proteins, that prevents the formation of structurally-defined complexes (Kay et al. 2000).

The effects of wPRP on solubility of individual milk proteins detected in supernatants after 15 min at 22 °C (Day 0) and 24 h at 4 °C (Day 1), were studied by CE. This experiment detected both the earliest interactions of wPRP peptides with milk proteins (15 min) and effects after extended storage 4 °C (Fig. 5a, b). The changes in concentrations of individual proteins in the supernatant suggested that initial interactions involved the major α S1 and α S1-P caseins (Fig. 5b) with solubility significantly elevated by ~ 30% after storage for 1 d. A similar trend was present but not significant at 15 min. wPRP also appeared to initially promote decreased solubility of β -lg but the effect was not significant after storage.

The preferential interaction of wPRP with α S1-Cn, the major casein protein accounting for ~ 40% of milk proteins by mass, can account for the bulk solubilisation effects by wPRP. Weakening of hydrophobic interactions at low temperature is known to cause release of β -Cn and κ -Cn from casein micelles (Kumosinski et al. 1993). However, the apparent effect of wPRP on α S-caseins supports that effects of wPRP on milk proteins are complimentary and additive to temperature-mediated relaxation of hydrophobic interactions, and further support observed synergistic effects of wPRP in combination with extended storage time, on increasing milk protein solubility.

The selective promotion of solubility of α S-caseins by wPRP can also be explained by the lower relative content of hydrophobic residues (Ala, Ile, Leu, Phe, Trp, Val) in α S-caseins (~ 30–32 mole %) compared with other caseins and major whey proteins (ranging from 33 to 40 mole %), and the consequently greater ease with which wPRP might disperse physical aggregates involving α S-caseins. By analogy that CLN interferes with β -sheet folding of A β 42 into fibrils, (Williamson, 1994; Kay et al. 2000), and that wPRP does likewise (Bharadwaj et al. 2012), it is proposed that wPRP selectively binds to β -sheet rich structures of milk proteins via proline-mediated interactions. Furthermore, proline binding to non-polar regions of proteins is known to reduce hydrophobicity resulting in partial dissociation of casein micelles (McMahon & Brown, 1984). In summary, we hypothesise that wPRP binds to the major milk protein, α S-caseins, via weak hydrophobic interactions probably β -sheet-rich domains, and acts like a 'strip bandage' that subsequently prevents hydrophobic associations, and possibly covalent reactivity, between milk proteins during and after heat treatment.

Potential applications of wPRP in dairy processing

In the context of dairy processing, potential applications for this novel wPRP product appear to be in stabilisation of protein concentrates or UHT products during shelf storage, to prevent or reverse progressive thickening and gelation. However, practical usefulness of wPRP for regulating protein aggregation in dairy products requires efficacy at the lowest ratio of wPRP to SMP. Heat stability of concentrated forms of milk proteins have been previously regulated by the addition of either phosphate or citrate salts. For concentrated forms of recombined milk products, phosphates are typically added at a concentration of 0.04 mol per kg skimmed milk solids (Augustin & Clarke, 1990). In comparison, inhibition of thermal aggregation of caseins (80 °C for 1 min), was achieved using 1 part hydrolysed lecithin to 18 parts casein (w/w) (Le et al. 2007). In the present study, the concentration of milk solids in the protein solubility assay was 10 mg/ml (or 3.7 mg/ml protein), representing approximately 1/10 of standard skimmed milk. In dose-response studies (Figs. 1b & 2), the ratio of wPRP to SMP solids required for moderate effect was approximately 1 part wPRP to 4 parts milk solids (w/w). However, the ongoing dispersion of physical aggregates by wPRP during storage could allow for lowering the initial concentration of wPRP required to achieve longer-term stabilisation, particularly with refrigeration. Further studies are needed to optimise the concentration ratio of wPRP required for stabilisation of protein solubility and to evaluate commercial feasibility of wPRP for this purpose.

Financial support from CSIRO Food and Nutritional Sciences and the Melbourne School of Land and Environment, University of Melbourne, and assistance with experimental procedures by Li Jiang Cheng and Katherine Robinson, are gratefully acknowledged.

References

- Anand K & Ward G 2005 Method of producing heat stable whey protein and products made there from. United States: US Patents Application No. US 2005/0163887 A1
- Anema SG 2008 The whey proteins in milk: thermal denaturation, physical interactions and effects on the functional properties of milk. In *Milk Proteins: From Expression to Food*, pp. 239–273 (Eds A Thompson, M Boland & H Singh). Amsterdam, Boston: Academic Press-Elsevier
- Anema SG & Li YM 2003 Association of denatured whey proteins with casein micelles in heated reconstituted skim milk and its effect on casein micelle size. *Journal of Dairy Research* **70** 73–83
- Augustin M-A & Clarke PT 1990 Effects of added salts on the heat stability of recombined concentrated milk. *Journal of Dairy Research* **57** 213–226
- Baldwin RL 1986 Temperature dependence of the hydrophobic interaction in protein folding. *Proceedings of the National Academy of Sciences* **83** 8069–8072
- Bennett L, Williams R, Ecroyd H, Liu YQ, Sudharmarajan S & Carver J 2009 Protective interactions of dairy peptides with fibril structures and relevance to Alzheimer's disease. *Australian Journal of Dairy Technology* **64** 117–121
- Bharadwaj P, Head R, Martins R, Raussens V, Sarroukh R, Jegasothy H, Waddington L & Bennett L 2012 Modulation of amyloid -b1-42 structure and toxicity by proline-rich whey peptides. *Food and Function*
- Boldogh I, Aguilera-Aguirre L, Bacsí A, Choudhury BK, Saavedra-Molina A & Kruzel M 2008 Colostrin decreases hypersensitivity and allergic

- responses to common allergens. *International Archives of Allergy and Immunology* **146** 298–306
- Chandrapala J, Augustin MA, McKinnon I & Udabage P** 2010 Effects of pH, calcium-complexing agents and milk solids concentration on formation of soluble protein aggregates in heated reconstituted skim milk. *International Dairy Journal* **20** 777–784
- Datta N & Deeth HC** 2001 Age gelation of UHT milk, a review. *Food and Bioprocesses Processing* **79** 197–210
- de Rham O & Chanton S** 1984 Role of ionic environment in insolubilization of whey protein during heat treatment of whey products. *Journal of Dairy Science* **67** 939–949
- Donato L & Guyomarch F** 2009 Formation and properties of the whey protein/kappa-casein complexes in heated skim milk – A review. *Dairy Science and Technology (EDP Sciences)* **89** 3–29
- Foegeding EA, Davis JP, Doucet D & McGuffey MK** 2002 Advances in modifying and understanding whey protein functionality. *Trends in Food Science and Technology* **13** 151–159
- Gao H, Havea P & Singh H** 2008 Whey product and process. USA: US Patents Application No. US 2008/0305235 A1
- Guyomarc'h F, Law AJR & Dalglish DG** 2003 Formation of soluble and micelle-bound protein aggregates in heated milk. *Journal of Agricultural and Food Chemistry* **51** 4652–4660
- Havn SS, Ipsen R, Nielsen PM & Lilbaek HM** 2006 Improving the foaming properties and heat stability of whey protein concentrates by phospholipase treatment. *Milchwissenschaft-Milk Science International* **61** 188–191
- Janusz M & Zablocka A** 2010 Colostral proline-rich polypeptides – immunoregulatory properties and prospects of therapeutic use in Alzheimers disease. *Current Alzheimer Research* **7** 323–333
- Janusz M, Staroscik K, Zimecki M, Wieczorek Z & Lisowski J** 1981 Chemical and physical characterisation of a proline-rich polypeptide from sheep colostrum. *Biochemical Journal* **199** 9–15
- Kanning M, Casella M & Olieman C** 1993 Milk and soy proteins analysis using capillary zone electrophoresis. *Liquid Chromatography-Gas Chromatography International* **6** 701–704
- Kay BK, Williamson MP & Sudol M** 2000 The importance of being proline: the interaction of proline-rich motifs in signaling proteins with their cognate domains. *FASEB Journal* **14** 231–241
- Kulmyrzaev A, Bryant C & McClements DJ** 2000 Influence of sucrose on the thermal denaturation, gelation, and emulsion stabilization of whey proteins. *Journal of Agricultural and Food Chemistry* **48** 1593–1597
- Kumosinski TF, Brown EM & Farrell HM** 1993 Three – dimensional molecular modeling of bovine caseins: a refined, energy-minimized kappa-casein structure. *Journal of Dairy Science* **76** 2507–2520
- Laleye LC, Jobe B & Wasesa AAH** 2008 Comparative study on heat stability and functionality of camel and bovine milk whey proteins. *Journal of Dairy Science* **91** 4527–4534
- Le TT, El-Bakry M, Neirynck N, Bogus M, Hoa HD & Van der Meeren P** 2007 Hydrophilic lecithins protect milk proteins against heat-induced aggregation. *Colloids and Surfaces B: Biointerfaces* **60** 167–173
- Le TT, Sabatino P, Heyman B, Kasinos M, Dinh HH, Dewettinck K, Martins J & Van der Meeren P** 2011 Improved heat stability by whey protein-surfactant interaction. *Food Hydrocolloids* **25** 594–603
- McCrae CH** 1999 Heat stability of milk emulsions: phospholipid-protein interactions. *International Dairy Journal* **9** 227–231
- McCrae CH & Muir DD** 1992 Heat stability of recombined milk: influence of lecithins on the heat coagulation time-pH profile. *Journal of Dairy Research* **59** 177–185
- McMahon DJ & Brown RJ** 1984 Composition, structure, and integrity of casein micelles: a review. *Journal of Dairy Science* **67** 499–512
- Morand M, Guyomarc'h F & Famelart MH** 2011 How to tailor heat-induced whey protein/kappa-casein complexes as a means to investigate the acid gelation of milk-a review. *Dairy Science and Technology* **91** 97–126
- Muir DD & Sweetsur AWM** 1977 Effect of urea on heat coagulation of caseinate complex in skim milk. *Journal of Dairy Research* **44** 249–257
- Muir DD & Sweetsur AWM** 1978 The effect of concentration on the heat stability of skim-milk. *Journal of Dairy Research* **45** 37–45
- O'Connell JE & Fox PF** 1999a Effect of extracts of oak (*Quercus petraea*) bark, oak leaves, aloe vera (Curacao aloe), coconut shell and wine on the colloidal stability of milk and concentrated milk. *Food Chemistry* **66** 93–96
- O'Connell JE & Fox PF** 1999b Effects of phenolic compounds on the heat stability of milk and concentrated milk. *Journal of Dairy Research* **66** 399–407
- O'Connell JE & Fox PF** 2003 Heat stability of milk. In *Encyclopedia of Dairy Sciences*, pp. 1321–1326 (Eds H Roginski, JW Fuquay & PF Fox). New York, Amsterdam: Academic Press
- Parris N** 1993 Aggregation of whey proteins in heated sweet whey. *Journal of Agricultural and Food Chemistry* **41** 460–464
- Parris N, Hollar CM, Hsieh A & Cockley KD** 1997 Thermal stability of whey protein concentrate mixtures: aggregate formation. *Journal of Dairy Science* **80** 19–28
- Raikos V** 2010 Effect of heat treatment on milk protein functionality at emulsion interfaces. A review. *Food Hydrocolloids* **24** 259–265
- Relkin P & Mulvihill DM** 1996 Thermal unfolding of β -lactoglobulin, α -lactalbumin, and bovine serum albumin. A thermodynamic approach. *Critical Reviews in Food Science and Nutrition* **36** 565–601
- Schuster D, Rajendran A, Hui SW, Nicotera T, Srikrishnan T & Kruzel ML** 2005 Protective effect of colostrin on neuroblastoma cell survival is due to reduced aggregation of beta-amyloid. *Neuropeptides* **39** 419–426
- Singh H** 2004 Heat stability of milk. *International Journal of Dairy Technology* **57** 111–119
- Thorn DC, Meehan S, Sunde M, Rekas A, Gras SL, MacPhee CE, Dobson CM, Wilson MR & Carver JA** 2005 Amyloid fibril formation by bovine milk kappa-casein and its inhibition by the molecular chaperones alpha(s-) and beta-casein. *Biochemistry* **44** 17027–17036
- Thorn DC, Ecroyd H, Sunde M, Poon S & Carver JA** 2008 Amyloid fibril formation by bovine milk alpha(s2)-casein occurs under physiological conditions yet is prevented by its natural counterpart, alpha(s1)-casein. *Biochemistry* **47** 3926–3936
- Thorn DC, Ecroyd H & Carver JA** 2009 The two-faced nature of milk casein proteins: amyloid fibril formation and chaperone-like activity. *Australian Journal of Dairy Technology* **64** 34–40
- Williams RPW, D'Ath L & Augustin MA** 2005 Production of calcium-fortified milk powders using soluble calcium salts. *Lait* **85** 369–381
- Williamson MP** 1994 The structure and function of proline-rich regions in proteins. *Biochemical journal* **297**(Pt 2) 249
- Xu RJ** 2001 Light scattering. In *Particle Characterisation: Light Scattering Methods*, pp. 56–100 Dordrecht, The Netherlands: Kluwer Academic Publishers
- Ye A** 2008 Complexation between milk proteins and polysaccharides via electrostatic interaction: principles and applications. A review. *International Journal of Food Science and Technology* **43** 406–415
- Yong YH & Forgeding EA** 2010 Caseins: utilizing molecular chaperone properties to control protein aggregation in foods. *Journal of Agricultural and Food Chemistry* **58** 685–693
- Yuksel Z & Erdem YK** 2005 The influence of main milk components on the hydrophobic interactions of milk protein system in the course of heat treatment. *Journal of Food Engineering* **67** 301–308
- Zimecki M** 2008 A proline-rich polypeptide from ovine colostrum: colostrin with immunomodulatory activity. In *Bioactive Components of Milk*, Vol. 606, pp. 241–250. Berlin: Springer-Verlag Berlin